

INFNCA-TH9709
 hep-ph/9706349
 June 1997

A GENERALIZATION OF THE BRODSKY-LEPAGE FORMALISM *

L. HOURA-YAOU, P. KESSLER, J. PARISI

*Laboratoire de Physique Corpusculaire, Collège de France
 11, Place Marcelin Berthelot, F-75231 Paris Cedex 05, France*

F. MURGIA

*Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari
 Casella Postale 170, I-09042 Monserrato (CA), Italy*

J. HANSSON

Department of Physics, Luleå University of Technology, S-95187 Luleå, Sweden

We present an approach that generalizes in a natural way the perturbative QCD formalism developed by Brodsky and Lepage for the study of exclusive hadronic processes to the case of $L \neq 0$ mesons. As an application of our approach we consider here the production of meson pairs, involving tensor and pseudotensor mesons, in photon-photon collisions.

In the last years a lot of theoretical work has been devoted to the study of hadronic exclusive processes at high transfer momentum, using factorization techniques and perturbative QCD (PQCD) ¹. While the PQCD formalism seems to be quite successful, it applies only to the case of hadrons with internal orbital angular momentum $L = 0$. Recently ² we have proposed an approach which generalizes the PQCD formalism to the case of the production and/or decay of mesons with a $(q\bar{q})$ valence Fock state and with any value of the orbital angular momentum L .

In this contribution we give a short, qualitative presentation of our approach, and discuss its application to the case of meson pair production in photon-photon collisions. Predictions for the production of pseudoscalar and vector meson pairs have been already given by Brodsky and Lepage ³. It is then interesting to extend this study to the production of other meson pairs, involving tensor and pseudotensor mesons.

Let us first recall how the helicity amplitudes for a generic exclusive hadronic process at high transfer momentum are evaluated in perturbative

*Talk delivered by F. Murgia at the PHOTON'97 Conference, Egmond aan Zee, The Netherlands, May 10-15, 1997.

QCD¹. Using factorization techniques, those amplitudes are expressed as a convolution among different contributions: *i*) A hard-scattering amplitude involving the valence partons of all participating hadrons, assumed to be, inside each hadron, in a collinear configuration; *ii*) Soft, nonperturbative distribution amplitudes (DA) for: *a*) finding the (collinear) valence partons in the incoming hadrons; *b*) the final partonic state to form the observed outgoing hadrons.

Notice that: *i*) Only leading Fock states are considered (i.e., $|q\bar{q}\rangle$ for mesons, $|qqq\rangle$ for baryons); *ii*) Valence parton masses are neglected (this, together with collinear configuration, forces the orbital angular momentum L of each hadron to be vanishing).

In our approach² we extend the PQCD formalism to the case of $L \neq 0$ mesons by combining it with a bound-state model of weakly bound valence quarks for the mesons involved. A general prescription can thus be given which relates the helicity amplitudes for the overall reaction (involving the observed mesons) to the partonic amplitudes (involving the collinear valence quarks of those mesons). This prescription implies the usual convolution of the partonic amplitude with the mesonic DA's, as discussed before. In addition, it involves also a convolution with the momentum-space wavefunction of the valence $q\bar{q}$ pair in each meson, given in the corresponding meson rest-frame. A Lorentz boost connects the rest frame of each meson with the center of mass frame of the initial particles; we assume that the mesons, as seen in this overall frame, are extreme-relativistic (this entails some simplifications in actual calculations). Another basic ingredient of our approach is the assumption that, for each meson, the momentum-space wavefunction of the valence $q\bar{q}$ pair is sharply peaked around zero, so that a series expansion in powers of the relative $q\bar{q}$ momentum can be made and only the leading terms have to be considered, in first approximation.

The following equation shows the resulting relationship between the overall and the partonic helicity amplitudes in the case of interest here, that is meson pair production in photon-photon collisions, $\gamma\gamma' \rightarrow QQ'$:

$$\begin{aligned}
M_{\lambda_\gamma \lambda_{\gamma'}}^{LSJ\Lambda, L'S'J'\Lambda'}(E, \Theta) &= f_L^* f_{L'}^* C_{0\Lambda\Lambda}^{LSJ} C_{0\Lambda'\Lambda'}^{L'S'J'} \\
&\times \lim_{\beta, \beta' \rightarrow 0} \frac{1}{\beta^L \beta'^{L'}} \int \frac{d(\cos \theta)}{2} d_{0,0}^L(\theta) \int \frac{d(\cos \theta')}{2} d_{0,0}^{L'}(\theta') \\
&\times \int \frac{\Phi_N^*(x) dx}{\sqrt{x(1-x)}} \int \frac{\Phi_N'^*(x') dx'}{\sqrt{x'(1-x')}} T_{\lambda_\gamma \lambda_{\gamma'}}^{S\Lambda, S'\Lambda'}(E, \Theta, \beta, \beta', \theta, \theta', x, x').
\end{aligned} \tag{1}$$

In this equation E and Θ are respectively the total energy and the scattering angle in the two-photon c.m. frame; L, S, J, Λ are the quantum numbers of meson Q (hereafter all corresponding labels with a prime refer to meson

Q'); f_L is the normalization constant related to the Q meson bound-state wavefunction; $\beta = 2|\mathbf{k}|/M_Q$, θ , refer to the relative $q\bar{q}$ momentum, $\mathbf{k}$, inside meson Q , in its rest frame; $\Phi_N(x)$ is the Q meson distribution amplitude; x is the fraction of Q meson longitudinal momentum carried by quark q ; finally, d -matrices and Clebsch-Gordan coefficients come from opportunely combining the $q\bar{q}$ spin and relative angular momentum states to fit the Q meson quantum numbers. Notice also that in the partonic amplitude T the helicities of each $q\bar{q}$ pair have already been combined to give the required spin and helicity for the corresponding meson.

Let us stress that Eq. (1) reduces exactly to the usual PQCD result when $L = 0$.

The partonic helicity amplitude T in Eq. (1) can be evaluated at leading order in perturbative QCD. The normalization constants f_L , $f_{L'}$ may be fixed by evaluating within the same theoretical approach the two-photon decay widths $\Gamma(Q \rightarrow \gamma\gamma)$ and comparing them with the corresponding available experimental results. In order to check the dependence of our results on the meson DA's appearing in Eq. (1) we have considered several indicative choices: *i)* For the pion, the asymptotic (ASY) DA, $\Phi_{N,\pi}^{ASY}(x) = 6x(1-x)$, and the Chernyak-Zhitnitsky (CZ) DA, $\Phi_{N,\pi}^{CZ}(x) = 30x(1-x)(2x-1)^2$; *ii)* For tensor and pseudotensor mesons the so-called nonrelativistic (NR) DA, $\Phi_N^{NR}(x) = \delta(x-1/2)$, and a generalized asymptotic (GASY) DA, $\Phi_N^L(x) \propto x^{L+1}(1-x)^{L+1}$, which is also required to get rid of possible divergences in the denominator of the partonic amplitudes (e.g., $\Phi_N^{L=1}(x) = 30x^2(1-x)^2$ for tensor mesons, $\Phi_N^{L=2}(x) = 140x^3(1-x)^3$ for pseudotensor mesons).

Let us now present some numerical results for the production of: *i)* Tensor meson pairs; *ii)* Pseudotensor meson pairs; *iii)* Pairs made by a pion and a pseudotensor meson. A more complete discussion can be found in Ref. 2; here we limit ourselves to present some indicative examples.

In Fig. 1 the scaling differential cross section $E^8 d\sigma/dt$, as a function of $\cos^2 \Theta$ is presented (Θ is the scattering angle in the two-photon c.m. frame). Results are given for a number of tensor meson pairs (using the GASY DA). For comparison, analogous results for pion pairs are also presented, using the asymptotic DA.

Tables 1 and 2 show, for all the meson pairs considered, the integrated cross sections (over the p_T of the mesons in the lab. frame, starting from a given minimum p_T) for the overall process $e^-e^+ \rightarrow e^-e^+QQ'$, in two kinematical conditions, corresponding roughly to those of LEP2 and of a possible B-factory. Again, several choices for the meson DA's are considered.

From the results of Fig. 1 and Tables 1,2 (see also Ref. 2) one can make the following conclusions: *i)* The results obtained do not depend strongly on the




Fig. 1: Differential cross section $E^8[d\sigma/dt]$ in $\text{nb}\times\text{GeV}^6$, as a function of $\cos^2 \Theta$, for the process $\gamma\gamma \rightarrow QQ'$ involving the production of tensor-meson pairs; the generalized asymptotic (GASY) DA was used for tensor mesons; for comparison, analogous curves for $\gamma\gamma \rightarrow \pi^+\pi^-$ and $\gamma\gamma \rightarrow \pi^0\pi^0$, using the asymptotic DA are also shown.

choice of the distribution amplitude for tensor and pseudotensor mesons; on the average, the GASY DA leads to results slightly (by a factor of 2 or 3) higher than the nonrelativistic one; *ii*) As expected, the charged channels give rise to significantly higher yields than the neutral ones, and are then the favorite candidates for experimental searches; *iii*) The production of the charged meson pairs here considered may become measurable with high-energy e^-e^+ colliders of the next generation, with integrated luminosities of about 10^{40} cm^{-2} .

Acknowledgments

This work has been partially supported by the EU program “Human Capital and Mobility” under contract CHRX-CT94-0450.

References

1. See, e.g., S.J. Brodsky and G.P. Lepage, in *Perturbative Quantum Chromodynamics*, ed. A.H. Mueller (World Scientific, Singapore, 1989).
2. L. Houra-Yaou, P. Kessler, J. Parisi, F. Murgia and J. Hansson, preprint hep-ph/9611337, *Z. Phys. C*, in press, 1997.
3. S.J. Brodsky and G.P. Lepage, *Phys. Rev. D* **24**, 1808 (1981).

Table 1: Integrated cross sections (in 10^{-40} cm^2) for the process $e^-e^+ \rightarrow e^-e^+QQ'$, at $\sqrt{s} = 200 \text{ GeV}$, $p_T > 1 \text{ GeV}$.

QQ'	$\sigma(e^-e^+ \rightarrow e^-e^+QQ') [10^{-40} \text{ cm}^2]$	
	NR	GASY
$f_2 f_2$	20.9	25.4
$a_2^0 a_2^0$	30.8	53.0
$f_2 a_2^0$	19.4	33.2
$f_2' f_2'$	0.6	1.1
$a_2^+ a_2^-$	291.0	1003.3
$\pi_2^0 \pi_2^0$	141.1	23.2
$\pi_2^+ \pi_2^-$	749.3	1418.2
$\pi_{CZ}^0 \pi_2^0$	116.6	263.0
$\pi_{ASY}^0 \pi_2^0$	35.2	104.1
$\pi_{CZ}^+ \pi_2^-$	4167.0	6562.6
$\pi_{ASY}^+ \pi_2^-$	1495.8	2368.5

Table 2: Same as table 1, but assuming: $\sqrt{s} = 10 \text{ GeV}$, $p_T > 1 \text{ GeV}$.

QQ'	$\sigma(e^-e^+ \rightarrow e^-e^+QQ') [10^{-40} \text{ cm}^2]$	
	NR	GASY
$f_2 f_2$	1.4	1.4
$a_2^0 a_2^0$	2.3	3.6
$f_2 a_2^0$	1.5	2.3
$f_2' f_2'$	0.04	0.06
$a_2^+ a_2^-$	20.1	68.6
$\pi_2^0 \pi_2^0$	10.0	1.6
$\pi_2^+ \pi_2^-$	41.6	72.4
$\pi_{CZ}^0 \pi_2^0$	12.0	24.5
$\pi_{ASY}^0 \pi_2^0$	3.0	8.8
$\pi_{CZ}^+ \pi_2^-$	341.1	536.7
$\pi_{ASY}^+ \pi_2^-$	122.0	193.2